

## Algebraic Topology – Homework 0

No due date

### Exercise 1.

- (a) Prove that, if  $(X, \mathcal{T})$  is a Hausdorff space and  $C \subseteq X$  is a compact subset, then  $C$  is closed.
- (b) Using (a), prove that if  $\pi: (P, \mathcal{T}_1) \longrightarrow (S, \mathcal{T}_2)$  is a continuous surjective map, with  $(P, \mathcal{T}_1)$  compact and  $(S, \mathcal{T}_2)$  Hausdorff, then  $\mathcal{T}_2$  coincides with the quotient topology.

The next exercise is a "Prove or disprove exercise" : it means that, if you "feel" that the assertion is true, you should prove it carefully ; if you "feel" it is false, you should find a contradiction. This type of exercises are harder but very instructive, as they improve intuition.

Here every subspace of  $(\mathbb{R}^n, \mathcal{E})$  is endowed with the subset topology, and  $\mathcal{E}$  denotes the Euclidean topology on  $\mathbb{R}^n$ ,

Let  $S^1 \subset \mathbb{R}^2$  be the 1-dimensional unit sphere, i.e. the unit circle

$$S^1 := \left\{ (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1 \right\},$$

and let  $\sim$  be the equivalence relation on  $S^1$  defined by :

$$(x_1, y_1) \sim (x_2, y_2) \iff (x_1, y_1) = \pm(x_2, y_2).$$

Then  $(\mathbb{R}P^1, \mathcal{T})$  is the topological space given by the equivalence classes of  $\sim$ , endowed with the quotient topology (with respect to the map that sends  $(x, y) \in S^1$  to  $[(x, y)] = \{(x, y), (-x, -y)\}$ ), and is called the **1-dimensional (real) projective space**. In other words, " $\mathbb{R}P^1$  is obtained from  $S^1$  by identifying antipodal points."

**Exercise 2.**

*Prove\* or disprove* the existence of the following homeomorphisms :

(a)  $\mathbb{R}P^1 \approx S^1$  ;

(b)  $S^1 \approx \infty$  (here " $\infty$ " is the subspace of  $\mathbb{R}^2$  given by two circles touching in one point, as in the infinity symbol) ;

(c)  $\mathbb{R}^1 \approx \mathbb{R}^m$ , with  $m \in \mathbb{Z}$ ,  $m > 1$  ;

(d)  $S^2 \setminus \{\mathbf{n}, \mathbf{s}\} \approx \mathcal{C}$ , where  $S^2$  is the unit sphere  $\{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ ,  $\mathbf{n} = (0, 0, 1)$ ,  $\mathbf{s} = (0, 0, -1)$  and  $\mathcal{C}$  is the cylinder given by  $\{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 = 1\}$ .

(e)  $\mathbb{R}^{n+1} \setminus \{0\} \approx S^n \times \mathbb{R}$ , where  $S^n$  is the  $n$ -dimensional unit sphere.

\* Here *Prove* means that you should find a homeomorphism *as explicitly as possible*.